

Asian Journal of Research in Biological and Pharmaceutical Sciences

Journal home page: www.ajrbps.com



ANTIFUNGAL ACTIVITY OF PETROLIUM AND ETHANOLIC EXTRACTS OF *MORINGA OLEIFERA* LEAVES AGAINST *CANDIDA ALBICANS* AND *ASPERGILLUS NIGER*

Kale Bhagwat*¹, Lonkar Amar¹, Jadhav Sourabh¹

¹*Department of Biochemistry, Pandit Deendayal Upadhyay Dental College and Hospital, Kegaon, Solapur (Maharashtra), India.

ABSTRACT

Moringa oleifera is well known medicinal plant. Its different parts are widely used for the treatment of different types of diseases since it has antibacterial and antifungal activity. The leaves are rich in iron, rhamnose and a unique group of compounds called glucosinolates and isothiocyanates. Other medical properties include antipyretic, antiepileptic, anti-inflammatory, antiulcerative, antihypertensive, cholesterol lowering, antioxidant, anti diabetic. The current investigation was undertaken to evaluate the antifungal activities by petroleum ether and ethanolic extracts of *Moringa oleifera* leaves against *Candida albicans* and *Aspergillus Niger*. From our study, it is found that as concentration of *Moringa* leaves extract increases the diameter of zone of inhibition found increased. The results were higher and effective for ethanolic extract than petroleum extract. The result were also found more effective against *Aspergillus Niger* than *Candida albicans*.

KEYWORDS

Moringa oleifera, *Aspergillus Niger*, Antifungal, Antibacterial and *Candida albicans*.

Author for Correspondence:

Kale Bhagwat,
Department of Biochemistry,
Pandit Deendayal Upadhyay Dental College and
Hospital,
Kegaon, Solapur (Maharashtra), India.

Email: bhagwatkale2644@rediffmail.com

INTRODUCTION

Moringa oleifera (Figure No.1) is well known, widely distributed natural species belongs to monogeneric family Moringaceae and order Violales (Table No.1)^{1,2}.

Moringa tree can grow well in the humid tropic or hot dry land with average height that ranges from 5 to 10 m. It can survive in harsh climatic condition including destitute soil without being much affected by drought³. These are true vascular plants contain xylem and phloem for conduction of water and nutrients respectively (Sub-kingdom: Tracheobionta). Its trunk is soft, white corky and

branches bearing a gummy bark. Each tripinnately compound leaves bear several small leaf legs^{4,5}. Moringa is flowering plant (Division: Angiospermae/Magnoliophyta). The flowers are fragrant, bisexual and pentapetalous. It contains five unequal, separate, thinly veined, yellowish-white petals. (Figure No.2) Flower is syncarpous and posses united carpels (Subclass: Dilleniidae). They grow on slender, hairy stalks in spreading or drooping later flower clusters. Leaves of these plants are feathery, pale green, compound and tripinnate. They are about 30–60 cm long and contains many small leaf lets^{6,7}. Each leaflet is about 1 to 2 cm long and 0.5 to 1.0 cm wide. Upper surface of leaf lets is smooth dark while lower is pale green. They are variable in size and shape (Figure No.3).

This plant is also known as drum stick tree or horse radish tree. It have many vernacular names such as kelor, marango, mlonge, moonga, mulangay, nebeday, saijhan, sajna or Ben oil tree^{8,9} *Moringa oleifera* is well known medicinal plant. Its different parts are widely used for the treatment of different types of diseases since is have antibacterial and antifungal activity. The leaves are rich in iron, rhamnase and a unique group of compounds called glucosinolates and isothiocyanates. Other medical proprieties include antipyretic, antiepileptic, anti-inflammatory, antiulcerative, antihypertensive, cholesterol lowering, antioxidant, anti diabetic^{10,11}. The current investigation was undertaken to evaluate the antifungal activities by petroleum ether and ethanolic extracts of *Moringa oleifera* leaves against *Candida albicans* and *Aspergillus Niger*.

MATERIAL AND METHODS

Collection of leaves

Naturally dried leaves were directly plucked from Moringa plant. The collection is carried out from different plants growing in area of Solapur city and surrounding villages. Plant leaves were further dried in an oven at 40°C for a total of three days and then finally ground to a fine powder by grinder. Extraction is carried out by using petroleum ether and ethyl alcohol¹²⁻¹⁶.

Preparation of leaf extract

20 gram of finely grinded powder of Moringa leaves were soaked in two different conical flasks

containing petroleum ether and ethyl alcohol respectively. The sample were shaken on rotary shaker at 200 rpm for 24 hrs. The extract were filtered using Whattman filter paper. Then solvent were evaporated to obtain dry extract using a rotary evaporator and were stored in refrigerator for antifungal analysis. Before testing five different concentration of extract were prepared for both solvents separately. This was carried by dissolving 100mg, 200mg, 300mg, 400mg and 500mg of extract in 1 ml solvent respectively⁽¹²⁻¹⁶⁾.

Antifungal activity

The antifungal activity of the *Moringa oleifera* leaf extracts was determined using agar well diffusion method by following the known procedure. Fungal strains were spread on the surface of agar plate aseptically by sterile cotton swab. When surface was little dried wells of 4mm diameter were punched with the help of sterile stainless steel boarer. 20µl of petroleum and ethanol extracts of Moringa leaves were loaded separately in wells in separate agar plates. Separate wells were used for different concentrations. Each plate further contained on well loaded with solvent act as negative control. While another well was loaded with 20µl antibiotic Nystatin (100mg/ml). The plates were incubated at 28°C for 72 hours and the antifungal activity was assessed by measuring the diameter of the zone of inhibition at the interval of 24hrs¹²⁻¹⁶.

RESULTS AND DISCUSSION

From our study, it is found that as concentration of Moringa leaves extract increases the diameter of zone of inhibition found increased. Against *Aspergillus Niger* zone of inhibition due to ethanolic extract of Moringa leaves was found increased from 7.1 mm to 17.8 mm as concentration increases from 100 to 500 mg per ml. While zone of inhibition due to petrolium extract of Moringa leaves was found increased from 4.7 mm to 13.2 mm as concentration increases from 100 to 500mg per ml. On other hand against *Candida albicans* zone of inhibition due to ethanolic extract of Moringa leaves was found increased from 5.1mm to 14.7mm as concentration increases from 100 to 500mg per ml. While zone of inhibition due to petrolium extract of Moringa leaves was found

increased from 2.9mm to 12.9mm as concentration increases from 100 to 500mg per ml.

Nystatin is a polyene antibiotic. Polyene antibiotic is a class of antimicrobial polyene compounds that target fungi. It is most commonly used for the treatment of fungal infections. It was used as a positive control against both *Aspergillus Niger* and *Candida albicans*.

The results were higher and effective for ethanolic extract than petroleum extract. The result were also found more effective against *Aspergillus Niger* (Graph No.1) than *Candida albicans* (Graph No.2).

Table No.1: Classification of *Moringa oleifera*

S.No	Botanical Classification	
1	Kingdom	Plantae
2	Sub-kingdom	Tracheobionta
3	Super-division	Spermatophyta
4	Division	Angiospermae/Magnoliophyta
5	Class	Magnoliopsida
6	Subclass	Dilleniidae
7	Order	Capparales
8	Family	Moringaceae
9	Genus	Moringa
10	Species	Oleifera




Figure No.1: *Moringa oleifera* tree




Figure No.2: *Moringa oleifera* flowers



Figure No.3: *Moringa oleifera* leaves



Graph No.1: Effect of Ethanolic extract (Red line) and Petroleum ether extract (Green line) of *Moringa oleifera* leaves against *Aspergillus Niger*



Graph No.2: Effect of Ethanolic extract (Red line) and Petroleum ether extract (Green line) of *Moringa oleifera* leaves against *Candida albicans*

CONCLUSION

From our study it comes to known that there are some alkaloids present in extract of leaves of *Moringa oleifera* which can be effectively used to treat candidosis and aspergillosis.

ACKNOWLEDGEMENT

The authors wish to express their sincere gratitude to Department of Biochemistry, Pandit Deendayal Upadhyay Dental College and Hospital, Kegaon, Solapur (Maharashtra), India for providing necessary facilities to carry out this research work.

CONFLICT OF INTEREST

We declare that we have no conflict of interest.

BIBLIOGRAPHY

1. Nadkarni A. Indian Materia Medica, Popular Prakashan: Bombay, 3rd Edition, 1976, 810-816.
2. Anwar F and Bhangar M. Analytical characterization of *Moringa oleifera* seed oil grown in temperate regions of Pakistan, *J. Agric. Food Chem*, 51(22), 2003, 6558-6563.
3. Anwar F *et al.* *Moringa oleifera*: a food plant with multiple medicinal uses, *Phytother. Res*, 21(1), 2007, 17-25.
4. Morton J F. The horseradish tree, *Moringa pterygosperma* (Moringaceae) - A boon to arid lands? *Econ. Bot*, 45(3), 1991, 318-333.
5. Gupta M *et al.* CNS activities of methanolic extract of *Moringa oleifera* root in mice, *Fitoterapia*, 70(3), 1999, 244-250.
6. Shukla S *et al.* Antifertility profile of the aqueous extract of *Moringa oleifera* roots, *Journal of Ethnopharmacology*, 22(1), 1998, 51-62.
7. Lurling M and Beekman W. Anti-cyano bacterial activity of *Moringa oleifera* seeds, *Journal Appl Phycol*, 22(4), 2010, 503-510.
8. Fahey J *et al.* The chemical diversity and distribution of glucosinolates and isothiocyanates among plants, *Phytochemistry*, 56(1), 2001, 5-51.
9. Bennett R *et al.* Profiling glucosinolates and phenolics in vegetative and reproductive tissues of the multi-purpose trees *Moringa oleifera* L. (Horseradish tree) and *Moringa stenopetala* L, *J Agric Food Chem*, 51(12), 2003, 3546-3553.
10. Pal S and Saha B. Studies on the antiulcer activity of M, *oleifera* leaf extract on gastric ulcer models in rats, *Phytother. Res*, 9(6), 1995, 463-465.
11. Bukar A and Oyeyi T. Antimicrobial profile of *Moringa oleifera* Lam. Extracts against some food-borne microorganisms, *Bayero Journal of Pure and Applied Sciences*, 3(1), 2010, 43-48.
12. Napoleon P *et al.* Isolation, analysis and identification of phytochemicals of antimicrobial activity of *Moringa oleifera* Lam, *Current Biotica*, 3(1), 2009, 33-37.
13. Nwosu M and Okafor J. Preliminary studies on the antifungal activities of some medicinal plants against *Basidiobolus* and some other pathogenic fungi, *Mycoses*, 38(5-6), 1995, 191-195.
14. Chuang P *et al.* *In vitro* antifungal activity of ethanolic extract of the leaves of *Moringa oleifera* against dermatophytes, *Bioresources Technology*, 98(1), 2007, 232-236.
15. Srinivasan D *et al.* Antimicrobial activity of certain Indian medicinal plants used in folkloric medicine, *Journal of Ethnopharmacology*, 74(3), 2001, 217-220.
16. Himal Paudel Chhetri *et al.* Phytochemical and antimicrobial evaluations of some medicinal plants of Nepal, *Kathmandu university journal of science, engineering and technology*, 1(5), 2008, 49-54.

Please cite this article in press as: Kale Bhagwat *et al.* Antifungal activity of petroleum and ethanolic extracts of *Moringa oleifera* leaves against *Candida albicans* and *Aspergillus Niger*, *Asian Journal of Research in Biological and Pharmaceutical Sciences*, 5(2), 2017, 86-90.